


What is claimed is:

1. A pharmaceutical composition for oral administration comprising a  $\delta$ -amino- $\gamma$ -hydroxy- $\omega$ -aryl-alkanoic acid amide renin inhibitor in an absorption enhancing carrier medium comprising:
  - (a) a lipophilic component;
  - (b) a high HLB surfactant; and
  - (c) a hydrophilic component;which composition upon admixing forms a stable microemulsion preconcentrate.
2. A pharmaceutical composition according to Claim 1, wherein the lipophilic component comprises a low HLB surfactant.
3. A pharmaceutical composition according to Claim 2, wherein the lipophilic component comprises a low HLB surfactant which is based on a medium or a long chain fatty acid, or a mixture of fatty acids thereof, and an oil which is a medium or a long chain fatty acid triglyceride, or a mixture of triglycerides thereof.
4. A pharmaceutical composition according to Claim 3, wherein the lipophilic component comprises a low HLB surfactant which is based on a medium chain fatty acid, or a mixture of fatty acids thereof, and an oil which is a medium chain fatty acid triglyceride, or a mixture of triglycerides thereof.
5. A pharmaceutical composition according to Claim 4, wherein the microemulsion preconcentrate is in the form of a water-in-oil microemulsion which upon administration or dilution with an aqueous medium spontaneously converts to an oil-in-water microemulsion.
6. A pharmaceutical composition according to Claim 4, wherein the  $\delta$ -amino- $\gamma$ -hydroxy- $\omega$ -aryl-alkanoic acid amide renin inhibitor has the formula



wherein  $R_1$  is  $C_{1-4}$ alkoxy- $C_{1-4}$ alkoxy or  $C_{1-4}$ alkoxy- $C_{1-4}$ alkyl;  $R_2$  is  $C_{1-4}$ alkyl or  $C_{1-4}$ alkoxy; and  $R_3$  and  $R_4$  are independently branched  $C_{3-4}$ alkyl; or a pharmaceutically acceptable salt thereof.

7. A pharmaceutical composition according to Claim 6, wherein the  $\delta$ -amino- $\gamma$ -hydroxy- $\omega$ -aryl-alkanoic acid amide renin inhibitor is a compound of formula (I) wherein  $R_1$  is 3-methoxypropoxy;  $R_2$  is methoxy; and  $R_3$  and  $R_4$  are isopropyl; or a pharmaceutically acceptable salt thereof.

8. A pharmaceutical composition according to Claim 7, wherein the  $\delta$ -amino- $\gamma$ -hydroxy- $\omega$ -aryl-alkanoic acid amide renin inhibitor is (2S,4S,5S,7S)-5-amino-4-hydroxy-2-isopropyl-7-[4-methoxy-3-(3-methoxy-propoxy)-benzyl]-8-methyl-nonanoic acid (2-carbamoyl-2-methyl-propyl)-amide hemifumarate.

9. A pharmaceutical composition according to Claim 8, wherein the microemulsion preconcentrate is in the form of a water-in-oil microemulsion which upon administration or dilution with an aqueous medium spontaneously converts to an oil-in-water microemulsion.

10. A pharmaceutical composition according to Claim 6, wherein the medium chain fatty acids of the lipophilic component have from 8 to 12 carbon atoms.

11. A pharmaceutical composition according to Claim 10, wherein the oil is selected from propylene glycol di-caprylate/caprate and glyceryl tri-caprylate/caprate.

12. A pharmaceutical composition according to Claim 6, wherein the low HLB surfactant has an HLB value ranging from about 2.5 to about 6.

13. A pharmaceutical composition according to Claim 6, wherein the high HLB surfactant has an HLB value ranging from about 13 to about 20.

14. A pharmaceutical composition according to Claim 13, wherein the high HLB surfactant is selected from polysorbate 80, macrogol-15 hydroxystearate, vitamin E-TPGS and polyoxyl 40 hydrogenated castor oil.

15. A pharmaceutical composition according to Claim 6, wherein the hydrophilic phase comprises PEG 300.

16. A pharmaceutical composition according to Claim 15, wherein the medium chain fatty acids of the lipophilic component have from 8 to 12 carbon atoms.
17. A pharmaceutical composition according to Claim 16, wherein the low HLB surfactant has an HLB value ranging from about 2.5 to about 6.
18. A pharmaceutical composition according to Claim 17, wherein the high HLB surfactant has an HLB value ranging from about 13 to about 20.
19. A pharmaceutical composition according to Claim 18, wherein the  $\delta$ -amino- $\gamma$ -hydroxy- $\omega$ -aryl-alkanoic acid amide renin inhibitor is a compound of formula (I) wherein  $R_1$  is 3-methoxypropoxy;  $R_2$  is methoxy; and  $R_3$  and  $R_4$  are isopropyl; or a pharmaceutically acceptable salt thereof.
20. A pharmaceutical composition according to Claim 19, wherein the oil is selected from propylene glycol di-caprylate/caprate and glyceryl tri-caprylate/caprate.
21. A pharmaceutical composition according to Claim 19, wherein the high HLB surfactant is selected from polysorbate 80, macrogol-15 hydroxystearate, vitamin E-TPGS and polyoxyl 40 hydrogenated castor oil.
22. A pharmaceutical composition according to Claim 19, wherein the  $\delta$ -amino- $\gamma$ -hydroxy- $\omega$ -aryl-alkanoic acid amide renin inhibitor is (2S,4S,5S,7S)-5-amino-4-hydroxy-2-isopropyl-7-[4-methoxy-3-(3-methoxy-propoxy)-benzyl]-8-methyl-nonanoic acid (2-carbamoyl-2-methyl-propyl)-amide hemifumarate.
23. A pharmaceutical composition according to Claim 22, wherein the oil is selected from propylene glycol di-caprylate/caprate and glyceryl tri-caprylate/caprate.
24. A pharmaceutical composition according to Claim 23, wherein the high HLB surfactant is selected from polysorbate 80, macrogol-15 hydroxystearate, vitamin E-TPGS and polyoxyl 40 hydrogenated castor oil.
25. A pharmaceutical composition according to Claim 24, wherein the microemulsion preconcentrate is in the form of a water-in-oil microemulsion which upon administration or dilution with an aqueous medium spontaneously converts to an oil-in-water microemulsion.

26. A method for the treatment of hypertension; congestive heart failure, cardiac hypertrophy, cardiac fibrosis, cardiomyopathy postinfarction, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, anxiety states and cognitive disorders which method comprises administering a therapeutically effective amount of a pharmaceutical composition according to Claim 1-24 or 25 to a patient in need thereof.

27. A pharmaceutical composition according to Claim 1-24 or 25, for use as medicament.

28. Use of a pharmaceutical composition according to Claim 1-24 or 25, for the manufacture of a medicament for the treatment of conditions associated with renin activity.